ERK 1/2 Polyclonal Antibody

Catalog Number:E-AB-31374 3 Publications

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity	Human,Mouse,Rat
Immunogen	Synthesized peptide derived from the C-terminal region of human ERK 1/2
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Formulation	PBS with 0.02% sodium azide, 0.5% BSA and 50% glycerol, pH7.4

Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:100-1:300
IF	1:50-1:200
ELISA	1:10000

Data



Western Blot analysis of K562 cells using ERK 1/2 Polyclonal Antibody at dilution of 1:2000.
Observed Mw:42,44kDa
Calculated Mw:43kDa



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using ERK 1/2 Polyclonal Antibody at dilution of 1:200.



Immunofluorescence analysis of Rat lung tissue using ERK 1/2 Polyclonal Antibody at dilution of 1:200.

Preparation & Storage

Storage	Store at -20℃. Avoid freeze / thaw cycles.

Background

Involved in both the initiation and regulation of meiosis, mitosis, and postmitotic functions in differentiated cells by

For Research Use Only

phosphorylating a number of transcription factors such as ELK1. Phosphorylates EIF4EBP1; required for initiation of translation. Phosphorylates microtubule-associated protein 2 (MAP2). Phosphorylates SPZ1 (By similarity). Phosphorylates heat shock factor protein 4 (HSF4) and ARHGEF2.Acts as a transcriptional repressor. Binds to a [GC]AAA[GC] consensus sequence. Repress the expression of interferon gamma-induced genes. Seems to bind to the promoter of CCL5, DMP1, IFIH1, IFITM1, IRF7, IRF9, LAMP3, OAS1, OAS2, OAS3 and STAT1. Transcriptional activity is independent of kinase activity.