CPSF4 Polyclonal Antibody

Catalog Number: E-AB-14891



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human, Mouse, Rat

Immunogen Recombinant protein of human CPSF4

Host Rabbit
Isotype IgG

Purification Affinity purification


Conjugation Unconjugated

Formulation PBS with 0.05% sodium azide, 50% glycerol, PH7.3

Applications Recommended Dilution

WB 1:500-1:2000

Data



Western Blot analysis of Hela and 293T cell using CPSF4 Polyclonal Antibody at dilution of 1:400

Calculated Mw:30kDa

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

Inhibition of the nuclear export of poly(A)-containing mRNAs caused by the influenza A virus NS1 protein requires its effector domain. The NS1 effector domain functionally interacts with the cellular 30 kDa subunit of cleavage and polyadenylation specific factor 4, an essential component of the 3' end processing machinery of cellular pre-mRNAs. In influenza virus-infected cells, the NS1 protein is physically associated with cleavage and polyadenylation specific factor 4, 30kD subunit. Binding of the NS1 protein to the 30 kDa protein in vitro prevents CPSF binding to the RNA substrate and inhibits 3' end cleavage and polyadenylation of host pre-mRNAs. Thus the NS1 protein selectively inhibits the nuclear export of cellular, and not viral, mRNAs. Multiple alternatively spliced transcript variants that encode different isoforms have been described for this gene.

For Research Use Only

Tel: 400-999-2100 Email: techsupport@elabscience.cn Web: www.elabscience.cn